Neuroscience Letters, 155 (1993) 237–239
Elsevier Scientific Publishers Ireland Ltd.

AUTHOR INDEX

Abe, K., see Aoki, M., 7
Aguirre, A., see Foret, J., 155
Alonso-Bedate, M., see Alonso-Gómez, A.L., 33
Alonso-Gómez, A.L., Alonso-Bedate, M. and Delgado, M.J., The inhibition by indoleamines (tryptamine and serotonin) of ocular serotonin-*N*-acetyltransferase from *Rana perezi* is temperature-dependent, 33
Aoki, M., Abe, K., Kawagoe, J., Nakamura, S. and Kogure, K., The preconditioned hippocampus accelerates HSP70 heat shock gene expression following transient ischemia in the gerbil, 7

Baimbridge, K.G., see Best, N., 1
Barajas-López, G., see González-Burgos, I., 99
Barker, J.L., see Valeyev, A.Y., 199
Bartlett, P.F., see Rosenfeld, J.V., 19
Beech, M.M., see Willoughby, J.J., 136
Benoit, O., see Foret, J., 155
Besson, M.-J., see Rogard, M., 81
Best, N., Mitchell, J., Baimbridge, K.G. and Wheal, H.V., Changes in parvalbumin-immunoreactive neurons in the rat hippocampus following a kainic acid lesion, 1
Bierer, L.M., see Gabriel, S.M., 116
Bloch, B., see Cadoret, M.-A., 92
Block, F., Schmitt, T. and Schwarz, M., Pentamidine, an inhibitor of spinal flexor reflexes in rats, is a potent *N*-methyl-D-aspartate (NMDA) antagonist in vivo, 208
Bruno, C., see Zadina, J.E., 220

Caboche, J., see Rogard, M., 81
Cadoret, M.-A., Jaber, M. and Bloch, B., Prenatal D_1, D_{1b} and D_3 dopamine receptor gene expression in the rat forebrain: detection by reverse polymerase chain reaction., 92
Candeo, P., see Manev, R.M., 73
Chen, H.I. and Liu, J.C., Localization of sympathetic postganglionic neurons innervating the femoral-saphenous vein in cats, 140
Clodoré, M., see Foret, J., 155
Couraud, F., see Grignon, S., 87
Cowburn, R.F., Vestling, M., Fowler, C.J., Ravid, R., Winblad, B. and O'Neill, C., Disrupted β_1-adrenoceptor—G protein coupling in the temporal cortex of patients with Alzheimer's disease, 163
Cox, R.D., see Naso, W.B., 125
Cruciani, R.A., see Valeyev, A.Y., 199

Davis, K.L., see Gabriel, S.M., 116
del Angel-Meza, A.R., see González-Burgos, I., 99
Delgado, M.J., see Alonso-Gómez, A.L., 33
der Linden, J.A.M.v., Joëls, M., Karst, H., Juta, A.J.A. and W.J.Wadman, , Bicuculline increases the intracellular calcium response of CA1 hippocampal neurons to synaptic stimulation, 230
Devasagayam, T.P.A., see Subramanian, M., 151
Domann, R., Hagemann, G., Kraemer, M., Freund, H.-J. and Witte, O.W., Electrophysiological changes in the surrounding brain tissue of photochemically induced cortical infarcts in the rat, 69
Donjo, T., see Miwa, T., 96

Eguchi, S., see Takahashi, M., 191

Fabbrizio, E., Leger, J., Leger, J.J. and Mornet, D., Dystrophin and dystrophin-related protein expression in *Torpedo marmorata* electric organ, 51
Fatome, M., see Martin, C., 77
Favaron, M., see Manev, R.M., 73
Feldman, A.G., The coactivation command for antagonist muscles involving Ib interneurons in mammalian motor control systems: an electrophysiologically testable model, 167
Feuerstein, G., see Jiang, Z.-G., 132
Finnman, U.-B., see Li, X.-M., 121
Fligny, I., see Willer, J.C., 212
Foret, J., Aguirre, A., Touitou, Y., Clodoré, M. and Benoit, O., Effect of morning bright light on body temperature, plasma cortisol and wrist motility measured during 24 hour of constant conditions, 155
Förster, E., Otten, U. and Frotscher, M., Developmental neurotrophin expression in slice cultures of rat hippocampus, 216
Förstermann, U., see Matsumoto, T., 61
Foster, R.G., see Provencio, I., 223
Fowler, C.J., see Cowburn, R.F., 163
Freund, H.-J., see Domann, R., 69
Frotscher, M., see Förster, E., 216
Funabashi, T., see Kimura, F., 183
Furukawa, M., see Miwa, T., 96
Fuxe, K., see Li, X.-M., 121

Gabellini, N., see Manev, R.M., 73
Gabriel, S.M., Bierer, L.M., Harotunian, V., Purohit, D.P., Perl, D.P. and Davis, K.L., Widespread deficits in somatostatin but not neuropeptide Y concentrations in Alzheimer's disease cerebral cortex, 116
Gajdusek, D.C., see Guiroy, D.C., 112
González-Burgos, I., del Angel-Meza, A.R. and Barajas-López, G., Morphometric characterization of the arcuate nucleus neurons of the rat. A Golgi study, 99
Grignon, S., Seagar, M.J. and Couraud, F., ω-Conotoxin sensitive calcium channels in cerebellar granule cells are not coupled to [^{3}H]glutamate release, 87
Guiroy, D.C., Marsh, R.F., Yanagihara, R. and Gajdusek, D.C., Immunolocalization of scrapie amyloid in non-congophilic, non-birefringent deposits in golden Syrian hamsters with experimental transmissible mink encephalopathy, 112

Hackler, L., see Zadina, J.E., 220
Hagemann, G., see Domann, R., 69
Harotunian, V., see Gabriel, S.M., 116
Hayakawa, M., see Itoh, Y., 144
Hedlund, P.B., see Li, X.-M., 121
Heimrich, *., see Olenik, C., 204
Hofmann, M.H. and Meyer, D.L., Visual-evoked responses in the salmon telencephalon change during smolt transformation, 234
Homma, S., see Takahashi, M., 191
Horn, D.A. and Latchman, D.S., Alternative splicing of the mRNA encoding the α subunits of the G_o GTP-binding protein during brain development and in neuronal cell lines, 57
Hossmann, K.-A., see Mies, G., 65

Iino, M., see Ozawa, S., 187

Ikeda, M., Morita, I., Murota, S.-i., Sekiguchi, F., Yuasa, T. and Miyatake, T., Cerebellar nitric oxide synthase activity is reduced in nervous and Purkinje cell degeneration mutants but not in climbing fiber-lesioned mice, 148
Ikeda, S.R., see Zhu, Y., 24
Itoh, Y., Yamada, M., Hayakawa, M., Otomo, E. and Miyatake, T., Subpial β/A4 peptide deposits are closely associated with amyloid angiopathy in the elderly, 144
Itoh, Y., see Sawada, M., 175

Jaber, M., see Cadoret, M.-A., 92
Jacobson, I., see Li, X., 42
Jiang, Z.-G., Yue, T.-L. and Feuerstein, G., Platelet-activating factor inhibits excitatory transmission in rat neostriatum, 132
Jinnai, K., see Kimura, F., 183
Joëls, M., see der Linden, J.A.M.v., 230
Julien, J.-F., see Rogard, M., 81
Juta, A.J.A., see der Linden, J.A.M.v., 230

Karst, H., see der Linden, J.A.M.v., 230
Kastin, A.J., see Zadina, J.E., 220
Kawagoe, J., see Aoki, M., 7
Kenigs, V., see Zadina, J.E., 220
Kimura, F., Jinnai, K. and Funabashi, T., A $GABA_B$-receptor mechanism is involved in the prolactin release in both male and female rats, 183
Kimura, Y., see Miwa, T., 96
Kogure, K., see Aoki, M., 7
Kohno, K., see Mies, G., 65
Kraemer, M., see Domann, R., 69
Kristensen, P., Suzdak, P.D. and Thomsen, C., Expression pattern and pharmacology of the rat type IV metabotropic glutamate receptor, 159
Krnjević, K., see Zhu, P.J., 128
Kuk, J.E., see Matsumoto, T., 61
Kuraishi, Y., see Ueda, M., 179
Kuroda, Y., Mikuni, M., Nomura, N. and Takahashi, K., Differential effect of subchronic dexamethasone treatment on serotonin-2 and β-adrenergic receptors in the rat cerebral cortex and hippocampus, 195

Lamas, G., see Willer, J.C., 212
Lange, G.D., see Valeyev, A.Y., 199
Latchman, D.S., see Horn, D.A., 57
Latchman, D.S., see Willoughby, J.J., 136
Leger, J., see Fabbrizio, E., 51
Leger, J.J., see Fabbrizio, E., 51
Li, X., Orwar, O., Persson, J., Sandberg, M. and Jacobson, I., γ-L-Glutamyl-L-glutamate is an endogenous dipeptide in the rat olfactory bulb which activates *N*-methyl-D-aspartate receptors, 42
Li, X.-M., Finnman, U.-B., von Euler, G., Hedlund, P.B. and Fuxe, K., Neuromedin N is a potent modulator of dopamine D_2 receptor agonist binding in rat neostriatal membranes, 121
Liu, J.C., see Chen, H.I., 140
Lobocki, C.A., see Pieper, D.R., 227
Lücke, T., see Neugebauer, V., 29

Manev, H., see Manev, R.M., 73
Manev, R.M., Favaron, M., Gabellini, N., Candeo, P. and Manev, H., Functional evidence for a L-AP3-sensitive metabotropic receptor different from glutamate metabotropic receptor mGluR1, 73
Marsh, R.F., see Guiroy, D.C., 112
Martin, C., Rubio, I. and Fatome, M., Early and transient effects of neutron irradiation on dopamine receptors in the adult rat brain, 77
Marunouchi, T., see Sawada, M., 175
Matsumoto, T., Nakane, M., Pollock, J.S., Kuk, J.E. and Förstermann, U., A correlation between soluble brain nitric oxide synthase and NADPH-diaphorase activity is only seen after exposure of the tissue to fixative, 61
McBryde, J.P., see Naso, W.B., 125
Metzinger, L., Passaquin, A.-C., Warter, J.-M. and Poindron, P., α-Methylprednisolone promotes skeletal myogenesis in dystrophin-deficient and control mouse cultures, 171
Meyer, D.K., see Olenik, C., 204
Meyer, D.L., see Hofmann, M.H., 234
Mies, G., Kohno, K. and Hossmann, K.-A., MK-801, a glutamate antagonist, lowers flow threshold for inhibition of protein synthesis after middle cerebral artery occlusion of rat, 65
Mikuni, M., see Kuroda, Y., 195
Mitchell, J., see Best, N., 1
Miwa, T., Moriizumi, T., Sakashita, H., Kimura, Y., Donjo, T. and Furukawa, M., Transection of the olfactory nerves induces expression of nerve growth factor receptor in mouse olfactory epithelium, 96
Miyamura, H., see Takahashi, M., 191
Miyatake, T., see Itoh, Y., 144
Miyatake, T., see Ikeda, M., 148
Moriizumi, T., see Miwa, T., 96
Morita, I., see Ikeda, M., 148
Mornet, D., see Fabbrizio, E., 51
Msghina, M. and Stjärne, L., Sympathetic transmitter release in rat tail artery and mouse vas deferens: facilitation and depression during high frequency stimulation, 37
Murota, S.-i., see Ikeda, M., 148

Nakamura, S., see Aoki, M., 7
Nakane, M., see Matsumoto, T., 61
Naso, W.B., Cox, R.D., McBryde, J.P. and Perot Jr., P.L., Rubrospinal neurons and retrograde transport of Fluoro-Gold in acute spinal cord injury — a dose–response curve, 125
Neugebauer, V., Lücke, T. and Schaible, H.-G., Differential effects of *N*-methyl-D-aspartate (NMDA) and non-NMDA receptor antagonists on the responses of rat spinal neurons with joint input, 29
Ninkina, N.N., see Willoughby, J.J., 136
Nomura, N., see Kuroda, Y., 195

O'Neill, C., see Cowburn, R.F., 163
Olenik, C., Heimrich, *. and Meyer, D.K., Expression of the cholecystokinin gene in organotypic slice cultures of immature rat somatosensory cortex, 204
Orwar, O., see Li, X., 42
Otomo, E., see Itoh, Y., 144
Otten, U., see Förster, E., 216
Otten, U., see Scully, J.L., 11
Ozawa, S. and Iino, M., Two distinct types of AMPA responses in cultured rat hippocampal neurons, 187

Pannicke, T., see Reichelt, W., 15
Passaquin, A.-C., see Metzinger, L., 171
Perl, D.P., see Gabriel, S.M., 116
Perot Jr., P.L., see Naso, W.B., 125
Persson, J., see Li, X., 42
Pieper, D.R. and Lobocki, C.A., Olfactory bulbectomy induces reproductive recrudescence in golden hamsters on short photoperiod, 227

Poindron, P., see Metzinger, L., 171
Pollock, J.S., see Matsumoto, T., 61
Provencio, I. and Foster, R.G., Vitamin A_2-based photopigments within the pineal gland of a fully terrestrial vertebrate, 223
Purohit, D.P., see Gabriel, S.M., 116
Pusphendran, C.K., see Subramanian, M., 151

Ravid, R., see Cowburn, R.F., 163
Reichelt, W. and Pannicke, T., Voltage-dependent K^+ currents in guinea pig Müller (glial) cells show different sensitivities to blockade by Ba^{2+}, 15
Richards, L.J., see Rosenfeld, J.V., 19
Rogard, M., Caboche, J., Julien, J.-F. and Besson, M.-J., The rat nucleus accumbens: two levels of complexity in the distribution of glutamic acid decarboxylase (67 kDa) and preproenkephalin messenger RNA, 81
Rosenfeld, J.V., Richards, L.J. and Bartlett, P.F., Mutant mouse cerebellum does not provide specific signals for the selective migration and development of transplanted Purkinje cells, 19
Rubio, I., see Martin, C., 77

Saji, M. and Volpe, B.T., Delayed histologic damage and neuron death in the substantia nigra reticulata following transient forebrain ischemia depends on the extent of initial striatal injury, 47
Sakashita, H., see Miwa, T., 96
Sandberg, M., see Li, X., 42
Satoh, M., see Ueda, M., 179
Sawada, M., Suzumura, A., Itoh, Y. and Marunouchi, T., Production of interleukin-5 by mouse astrocytes and microglia in culture, 175
Schaible, H.-G., see Neugebauer, V., 29
Schmitt, T., see Block, F., 208
Schwarz, M., see Block, F., 208
Scully, J.L. and Otten, U., Glucocorticoid modulation of neurotrophin expression in immortalized mouse hippocampal neurons, 11
Seagar, M.J., see Grignon, S., 87
Sekiguchi, F., see Ikeda, M., 148
Smallwood, V.S., see Valeyev, A.Y., 199
Sokoloff, A.J., Topographic segregation of genioglossus motoneurons in the neonatal rat, 102
Soudant, J., see Willer, J.C., 212
Stjärne, L., see Msghina, M., 37
Subramanian, M., Pusphendran, C.K., Tarachand, U. and Devasagayam, T.P.A., Gestation confers temporary resistance to peroxidation in the maternal rat brain, 151
Suzdak, P.D., see Kristensen, P., 159
Suzumura, A., see Sawada, M., 175

Takahashi, K., see Kuroda, Y., 195
Takahashi, M., Miyamura, H., Eguchi, S. and Homma, S., The effect of bupivacaine hydrochloride on skeletal muscle fiber type transformation by low frequency electrical stimulation, 191
Tam, S.W., see Zaczek, R., 107
Tarachand, U., see Subramanian, M., 151
Thomsen, C., see Kristensen, P., 159
Tinker, W.J., see Zaczek, R., 107
Touitou, Y., see Foret, J., 155

Ueda, M., Kuraishi, Y. and Satoh, M., Detection of capsaicin-evoked release of glutamate from spinal dorsal horn slices of rat with on-line monitoring system, 179

Valeyev, A.Y., Cruciani, R.A., Lange, G.D., Smallwood, V.S. and Barker, J.L., Cl^- channels are randomly activated by continuous GABA secretion in cultured embryonic rat hippocampal neurons, 199
Vestling, M., see Cowburn, R.F., 163
Volpe, B.T., see Saji, M., 47
von Euler, G., see Li, X.-M., 121

Warter, J.-M., see Metzinger, L., 171
Wheal, H.V., see Best, N., 1
Willer, J.C., Lamas, G., Fligny, I. and Soudant, J., Hypoglossal-facial anastomosis alters excitability of hypoglossal motoneurones in man, 212
Willoughby, J.J., Ninkina, N.N., Beech, M.M., Latchman, D.S. and Wood, J.N., Molecular cloning of a human neuronal nicotinic acetylcholine receptor β3-like subunit, 136
Winblad, B., see Cowburn, R.F., 163
Witte, O.W., see Domann, R., 69
W.J.Wadman, , see der Linden, J.A.M.v., 230
Wood, J.N., see Willoughby, J.J., 136

Yamada, M., see Itoh, Y., 144
Yanagihara, R., see Guiroy, D.C., 112
Yuasa, T., see Ikeda, M., 148
Yue, T.-L., see Jiang, Z.-G., 132

Zaczek, R., Tinker, W.J. and Tam, S.W., Unique properties of norepinephrine release from terminals arising from the locus coeruleus: high potassium sensitivity and lack of linopirdine (DuP 996) enhancement, 107
Zadina, J.E., Kastin, A.J., Kenigs, V., Bruno, C. and Hackler, L., Prolonged analgesia after intracerebroventricular Tyr-W-MIF-1 (Tyr-Pro-Trp-Gly-NH_2), 220
Zhu, P.J. and Krnjević, K., Adenosine release is a major cause of failure of synaptic transmission during hypoglycaemia in rat hippocampal slices, 128
Zhu, Y. and Ikeda, S.R., 2,3-Butanedione monoxime blockade of Ca^{2+} currents in adult rat sympathetic neurons does not involve 'chemical phosphatase' activity, 24